## **RELATION BETWEEN NERVE FIBER TYPE LESION AND NEURODYNAMIC TESTS IN PATIENTS WITH LUMBOSACRAL RADICULAR PAIN**



Pesesse P.<sup>1,2</sup>, Demoulin C.<sup>1,2</sup>, Stormacq G.<sup>1,2</sup>, Vanderthommen M.<sup>1,2</sup>

<sup>1</sup> Department of Sport and Rehabilitation Sciences, University of Liege, Liege, Belgium <sup>2</sup> Department of Physical Medicine and Rehabilitation, Liege University Hospital Center, Liege, Belgium



Lumbosacral radicular pain (LRP) is an important health care burden with a prevalence ranging from 1.2 to 43%. Neurodynamic tests are frequently used for the physical examination of nerve related pain. Baselgia and al. reported that negative neurodynamic tests are correlated with pronounced small nerve fiber sensory deficit in patients suffering from median nerve compression (1). Nonetheless, such a correlation has never been studied yet regarding LRP.



## II. METHODS

The study consisted of a single assessment session. In order to be included patients had to suffer from LRP associated with a positive slump test or a nerve dysfunction pointed out by means muscle manual testing (nerve Of root corsesponding myotome), tendon reflex examination (patellar and ankle) and/or clinical sensory testing (Figure 2). Six clinical test developed to evaluate small nerve fiber function were included in the clinical sensory testing.





Figure 2 : Clinical sensor testing, from Zhu, G. C. et al. Concurrent validity of a low-cost and time-efficient clinical sensory test battery to evaluate somatosensory dysfunction. Eur. J. Pain (United Kingdom) 23, 1826–1838

(2019)

## **III. RESULTS**

Among the 8 patients recruited, 6 had a positive slump test and 7 presented at least one positive sensory test for the small nerve fibers (Figure 1). Patients presenting with a negative slump test did not have a greater small nerve fibers dysfunction than patients with a positive neurodynamic test. Furthermore no specific small nerve fibers type dysfunction was linked to a negative neurodynamic test.

	Positive small nerves fibers test	Slump test results
Patient 1	0/6	Positive
Patient 2	1/6	Positive
Patient 3	1/6	Positive
Patient 4	2/6	Negative
Patient 5	3/6	Positive
Patient 6	3/6	Negative
Patient 7	5/6	Positive
Patient 8	5/6	Positive

Figure 3 : Results of small nerve fibers tests and neurodynamic test in the 8 participants

## **IV. CONCLUSION**

In this pilot study with patients suffering from LRP, negative neurodynamic tests did not appear to be associated with more small nerve fiber dysfunctions. However, considering the small sample size, these results need to be confirmed in further studies



- Baselgia, L. T., et al. Negative neurodynamic tests do not exclude neural dysfunction in patients with entrapment neuropathies. Arch. Phys. Med. Rehabil. 98, 480–486 (2017).
- Konstantinou K, Dunn KM. Sciatica: review of epidemiological studies and prevalence estimates. Spine (Phila Pa 1976). 2008;33(22):2464–72.

